

INDEPENDENT SETS IN EDGE-CLIQUE GRAPHS

Ching-Hao Liu, Ton Kloks, and Sheung-Hung Poon

Department of Computer Science
National Tsing Hua University, Taiwan
chinghao.liu@gmail.com spoon@cs.nthu.edu.tw

Abstract. We first show that the independent set problem on edge-clique graphs of cographs can be solved in $O(n^2)$ time. We then show that the independent set problem on edge-clique graphs of graphs without odd wheels is NP-complete. We present a PTAS for planar graphs and show that the problem is polynomial for planar graphs without triangle separators. Lastly, we show that edge-clique graphs of cocktail party graphs have unbounded rankwidth.

1 Introduction

Let $G = (V, E)$ be an undirected graph with vertex set V and edge set E . A clique is a complete subgraph of G .

Definition 1. *The edge-clique graph $K_e(G)$ of a graph G has the edges of G as its vertices and two vertices of $K_e(G)$ are adjacent when the corresponding edges in G are contained in a clique.*

Edge-clique graphs were introduced and studied by Albertson and Collins [1]. Two characterizations of edge-clique graphs have been presented by [10,12], but the problem whether there is any polynomial-time recognition algorithm for edge-clique graphs remains open. Some results of edge-clique graphs can be found in [1,8,9,10,11,12,26,29,36,38,39]. Notice that edge-clique graphs were implicitly first used by Kou et al [28]. In the following we introduce the independent set problem.

An independent set in a graph G is a nonempty set of vertices A such that there is no edge between any pair of vertices in A . The independent set problem asks the maximal cardinality of independent sets, which is called independent set number and denoted by $\alpha(G)$. In general graphs, this problem is known to be NP-hard and it remains NP-hard even for triangle-free graphs [35] and planar graphs of degree at most three [18]. In some classes of graphs, for example claw-free graphs [41] and perfect graphs [20], this problem can be solved in polynomial time.

A subset A of edges in a graph G is independent if it induces an independent set in $K_e(G)$, that is, no two edges of A are contained in a clique of G . We denote

the maximal cardinality of independent sets of edges of G by $\alpha'(G) = \alpha(K_e(G))$. Notice that, for any graph G , the clique number of its edge-clique graph satisfies

$$\omega(K_e(G)) = \binom{\omega(G)}{2}.$$

For the independent set number $\alpha(K_e(G))$ there is no such relation. For example, when G has no triangles then $K_e(G)$ is an independent set and the independent set problem in triangle-free graphs is NP-complete.

There is another problem relating to independent sets in edge-clique graphs, which is called the edge-clique cover problem. An edge-clique covering of G is a family of complete subgraphs such that each edge of G is in at least one member of the family. The edge-clique cover problem asks the minimal cardinality of such a family, which is called the edge-clique covering number and denoted by $\theta_e(G)$.

The problem of deciding if $\theta_e(G) \leq k$, for a given natural number k , is NP-complete [23,28,33]. The problem remains NP-complete when restricted to graphs with maximum degree at most six [24]. Hoover [24] gave a polynomial time algorithm for graphs with maximum degree at most five. For graphs with maximum degree less than five, this was already done by Pullman [37]. Also for linegraphs the problem can be solved in polynomial time [33,37]. In [28] it was shown that approximating the clique covering number within a constant factor smaller than two remains NP-complete. The independence number of $K_e(G)$ equals $\theta_e(G)$ for graphs G that are chordal. For interval graphs the edge-clique covering number $\theta_e(G)$ equals the number of maximal cliques [40].

We organized this paper as follows. In Section 2 and Section 3 we show that the independent set problem on edge-clique graphs is NP-hard in general and remains so for edge-clique graphs of graphs without odd wheels. We show in Sections 2.1 and 2.2 that the problem can be solved in $O(n^2)$ time for cographs and in polynomial time for distance-hereditary graphs. In Section 4, we present a PTAS for planar graphs and show that the problem is polynomial for planar graphs without triangle separator.

Notice that cographs have rankwidth one. If edge-clique graphs of cographs G would have bounded rankwidth, then computing $\alpha'(G)$ would be easy, because computing the independence number is polynomial for graphs of bounded rankwidth. However, we show in Section 5 that this is not the case, not even for edge-clique graphs of cocktail parties. Finally, we conclude in Section 6.

2 Algorithms for cographs

In this section we begin with a NP-hardness proof for the computation of $\alpha'(G)$ for arbitrary graphs G . Then we provide algorithms for the computation of independence number of edge-clique graphs of cographs and distance-hereditary graphs.

The following lemma shows that the independent set problem in $K_e(G)$ can be reduced to the independent set problem in G .

Lemma 1. *The computation of $\alpha'(G)$ for arbitrary graphs G is NP-hard.*

Proof. Let G be an arbitrary graph. Construct a graph H as follows. At every edge in G add two simplicial vertices¹, both adjacent to the two endvertices of the edge. Add one extra vertex x adjacent to all vertices of G . Let H be the graph constructed in this way.

Notice that a maximum set of independent edges does not contain any edge of G since it would be better to replace such an edge by two edges incident with the two simplicial vertices at this edge. Also notice that a set of independent edges incident with x corresponds with an independent set of vertices in G . Hence

$$\alpha'(H) = 2m + \alpha(G),$$

where m is the number of edges of G . □

2.1 Cographs

A cograph is a graph without induced P_4 . It is well-known that a graph is a cograph if and only if every induced subgraph with at least two vertices is either a join or a union of two smaller cographs. It follows that a cograph G has a decomposition tree (T, f) where T is a rooted binary tree and f is a bijection from the vertices of G to the leaves of T . Each internal node of T , including the root, is labeled as $\otimes$ or $\oplus$. The $\otimes$ -node joins the two subgraphs mapped to the left and right subtree. The $\oplus$ unions the two subgraphs. When G is a cograph then a decomposition tree as described above can be obtained in linear time [13].

Lemma 2. *Let G be a cograph. Assume that G is the join of two smaller cographs G_1 and G_2 . Then any edge in G_1 is adjacent in $K_e(G)$ to any edge in G_2 .*

For a vertex x , let $d'(x)$ be the independence number of the subgraph of G induced by $N(x)$, that is,

$$d'(x) = \alpha(G[N(x)]). \quad (1)$$

Lemma 3. *Let G be a cograph. Then*

$$\alpha'(G) = \max \left\{ \sum_{x \in W} d'(x) \mid W \text{ is an independent set in } G \right\}. \quad (2)$$

Proof. Cographs are characterized by the fact that every induced subgraph has a twin. Let x be a false twin of a vertex y in G . Let A be a maximum independent set of edges in G . Let $A(x)$ and $A(y)$ be the sets of edges in A that are incident with x and y . Assume that $|A(x)| \geq |A(y)|$. Let $\Omega(x)$ be the set of endvertices in $N(x)$ of edges in $A(x)$. then we may replace the set $A(y)$ with the set

$$\{ \{y, z\} \mid z \in \Omega(x) \}.$$

¹ An additional argument would show that adding one simplicial per edge is sufficient to prove NP-hardness.

The cardinality of the new set is at least as large as $|A|$. Notice that, for any maximal independent set Q in G , either $\{x, y\} \subseteq Q$ or $\{x, y\} \cap Q = \emptyset$. By induction on the number of vertices in G , Equation (2) is valid.

Let x be a true twin of a vertex y in G . Let A be a maximum independent set of edges in G and let $A(x)$ and $A(y)$ be the sets of edges in A that are incident with x and y . If $\{x, y\} \in A$ then $A(x) = A(y) = \{\{x, y\}\}$.

Now assume that $\{x, y\} \notin A$. Endvertices in $N(x)$ of edges in $A(x)$ and $A(y)$ are not adjacent nor do they coincide. Replace $A(x)$ with

$$\{ \{x, z\} \mid \{x, z\} \in A(x) \text{ or } \{y, z\} \in A(y) \}$$

and set $A(y) = \emptyset$. Then the new set of edges is independent and has the same cardinality as A .

Let Q be an independent set in G . At most one of x and y is in Q . By induction on the number of vertices in G , the validity of Equation (2) is easily checked. $\square$

Remark 1. Notice that the righthand side of Equation (2) is a lowerbound for $\alpha'(G)$ for any graph G . In this paper we find that there is an equality for some classes of graphs. Notice that, for example, equality does not hold for C_5 .

Theorem 1. *When G is a cograph then $\alpha'(G)$ satisfies Equation (2). This value can be computed in $O(n^2)$ time.*

Proof. We proved that $\alpha'(G)$ satisfies Equation (2) in Lemma 3. Here we show that $\alpha'(G)$ can be computed in $O(n^2)$ time.

The algorithm first computes a decomposition tree (T, f) for G in linear time. For each node p of T let G_p be the subgraph induced by the set of vertices that are mapped to leaves in the subtree rooted at p . Notice that the independence number of each G_p can be computed in linear time.

Compute $d'(x)$ for $x \in V$ as follows. Assume G is the union of $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$. For $i \in \{1, 2\}$, let $d'_i(x) = \alpha(G_i[N(x)])$ for vertices x in G_i . Then

$$d'(x) = \begin{cases} d'_1(x) & \text{if } x \in V_1 \\ d'_2(x) & \text{if } x \in V_2. \end{cases} \quad (3)$$

Assume that G is the join of G_1 and G_2 . Then

$$d'(x) = \begin{cases} \max \{ d'_1(x), \alpha(G_2) \} & \text{if } x \in V_1 \\ \max \{ d'_2(x), \alpha(G_1) \} & \text{if } x \in V_2. \end{cases} \quad (4)$$

Evaluate $\alpha'(G)$ as follows. Assume that G is the union of G_1 and G_2 . Then

$$\alpha'(G) = \alpha'(G_1) + \alpha'(G_2). \quad (5)$$

Assume that G is the join of G_1 and G_2 . For $i \in \{1, 2\}$, let

$$\alpha'_i = \max \left\{ \sum_{x \in W} d'(x) \mid W \text{ is an independent set in } G_i \right\}. \quad (6)$$

Then

$$\alpha'(G) = \max \{ \alpha'_1, \alpha'_2 \}. \quad (7)$$

This completes the proof. $\square$

Remark 2. For any graph G whose edge-clique graph is perfect the intersection number of G equals the fractional intersection number of G [26, Theorem 4.1]. It is easy to see that for cographs G , $K_e(G)$ is not necessarily perfect. For example, when G is the join of P_3 and C_4 , then $K_e(G)$ contain a C_5 as an induced subgraph.²

2.2 Distance-hereditary graphs

A graph G is distance-hereditary if the distance between any two nonadjacent vertices, in any connected induced subgraph of G , is the same as their distance in the G [25]. Bandelt and Mulder obtained the following characterization of distance-hereditary graphs.

Lemma 4 ([5]). *A graph is distance-hereditary if and only if every induced subgraph has an isolated vertex, a pendant vertex or a twin.*

The papers [5] and [25] also contain characterizations of distance-hereditary graphs in terms of forbidden induced subgraphs.

Let G be distance-hereditary. Then $\alpha'(G)$ does not necessarily satisfy Equation (2). A counterexample is as follows. Take two 4-wheels and join the two centers by an edge. The maximum edge-independent set consists of the edges of the two 4-cycles and the edge joining the two centers. However, there is no independent set such that (2) gives this value.

3 NP-completeness for graphs without odd wheels

A wheel W_n is a graph consisting of a cycle C_n and one additional vertex adjacent to all vertices in the cycle. The universal vertex of W_n is called the hub. It is unique unless $W_n = K_4$. The edges incident with the hub are called the spokes of the wheel. The cycle is called the rim of the wheel. A wheel is odd if the number of vertices in the cycle is odd.

² Let $P_3 = \{1, 2, 3\}$ and $C_4 = \{4, 5, 6, 7\}$. There is a C_5 with vertices

$$\{\{1, 2\}, \{3, 4\}, \{3, 7\}, \{4, 5\}, \{6, 7\}\}.$$

Lakshmanan, Bujtás and Tuza investigate the class of graphs without odd wheels in [30]. They prove that Tuza’s conjecture holds true for this class of graphs (see also [22]). Notice that a graph G has no odd wheel if and only if every neighborhood in G induces a bipartite graph. It follows that $\omega(G) \leq 3$. Obviously, the class of graphs without odd wheels is closed under taking subgraphs. Notice that, when G has no odd wheel then every neighborhood in $K_e(G)$ is either empty or a matching. Furthermore, it is easy to see that $K_e(G)$ contains no diamond (every edge is in exactly one triangle), no C_5 and no odd antihole.

For graphs G without odd wheels $K_e(G)$ coincides with the anti-Gallai graphs introduced by Le [32], since $\omega(G) \leq 3$. For general anti-Gallai graphs the computation of the clique number and chromatic number are NP-complete.

Let us mention that the recognition of anti-Gallai graphs is NP-complete. Even when each edge in G is in exactly one triangle, the problem to decide if G is an anti-Gallai graph is NP-complete [3, Corollary 5.2]. The recognition of edge-clique graphs of graphs without odd wheels is, as far as we know, open. Let us also mention that the edge-clique graphs of graphs without odd wheels are clique graphs [9]. The recognition of clique graphs of general graphs is NP-complete [2].

Theorem 2. *The computation of $\alpha'(G)$ is NP-complete for graphs G without odd wheels.*

Proof. We reduce 3-SAT to the vertex cover problem in edge-clique graphs of graphs without odd wheels.

Let $H \simeq cp(3)$, ie, $L(K_4)$. See Fig. 1(A). Let S be a 3-sun as depicted in Fig. 1(B). The graph H is obtained from S by adding three edges between pairs of vertices of degree two in S .³ Call the three vertices of degree four in S , the ‘inner triangle’ of H and call the remaining three vertices of H the ‘outer triangle.’

Notice that H has 3 maximum independent sets of edges. Each maximum independent set of edges is an induced C_4 consisting of one edge from the inner triangle, one edge from the outer triangle, and two edges between the two triangles. The three independent sets partition the edges of H .

The six edges of H between the inner and outer triangle form a 6-cycle in $K_e(H)$. Let F denote this set of edges in H .

For each clause $(x_i \vee x_j \vee x_k)$ take one copy of H . Take an independent set of three edges contained in F and label these with x_i , x_j and x_k .

For each variable x take a triangle. Label one edge of the triangle with the literal x and one other edge of the triangle with its negation $\bar{x}$. Then add a simplicial vertex to the unlabeled edge of the triangle.

Construct links between variable gadgets and clause gadgets as follows. Let $(x_i \vee x_j \vee x_k)$ be a clause. There are four steps. First add a 2-chain pv_1q

³ In [31, Theorem 14] the authors prove that every maximal clique in $K_e(G)$ contains a simplicial vertex if and only if G does not contain, as an induced subgraph, K_4 nor a 3-sun with 0, 1, 2 or 3 edges connecting the vertices of degree two.




Fig. 1: This figure shows H , S and $K_e(H)$. In H , vertices 2, 4, 5 induce the inner triangle and the remaining three vertices induce the outer triangle. The three colors for the edges of H indicate the partition of three maximum independent sets of edges.




Fig. 2: This figure shows the construction of a link between a variable gadget and a clause gadget.

from an endpoint p of the edge labeled x_i in the variable gadget to one endpoint q of the edge x_i in the clause gadget. See Fig. 2. Similarly, add a 3-chain $p'v_2v_3q'$ between the other two endpoints, where p' is in the variable gadget and q' is in the clause gadget. Then add four edges pv_2 , v_2v_1 , v_1v_3 and v_3q to let the rectangle $pqq'p'$ become a triangle strip with five triangles. Finally, add a simplicial vertex to each of the edges in the 2-chain and the 3-chain. We construct links for the other two literals in the clause in the same manner.

Let G be the graph constructed in this manner. Consider the triangles T which are the edge-clique graphs of the triangles in G containing the newly-added simplicial vertices. Notice that simplicial vertices of a graph may be removed without changing the complexity of the vertex cover problem and so the vertices of T can be removed without changing the complexity. Let K be the graph obtained from $K_e(G)$ by removing the edges of these triangles.

Let L be the number of variables, let M be the number of clauses in the 3-SAT formula. Assume that there is a satisfying assignment. Then choose the vertices in K corresponding to literals that are TRUE in the vertex cover. The variable gadgets need L vertices and the links need $6M$ vertices in the vertex cover. Since this assignment is satisfying, we need at most $8M$ vertices to cover the remaining edges in the clause structures, since the outgoing edge from each literal which is TRUE is covered. Thus there is a vertex cover of $K_e(G)$ with $L + 14M$ vertices.

Assume that $K_e(G)$ has a vertex cover with $L + 14M$ vertices. At least L vertices in K are covering the edges in the variable gadgets and at least $6M$ vertices in K are covering the edges in the links. The other $8M$ vertices of K are covering the edges in the clause gadgets. Take the literals of the variable gadgets that are in the vertex cover as an assignment for the formula. Each clause gadget must have one literal vertex of which the outgoing edge is covered. Therefore, the assignment is satisfying.

This proves the theorem. $\square$

4 Algorithms for planar graphs

In this section we obtain a PTAS for a maximum independent set of edges in planar graphs.

Lemma 5. *Let $k \in \mathbb{N}$. There exists a linear-time algorithm which computes a maximum independent set of edges for graphs of treewidth at most k .*

Proof. Notice that the problem can be formulated in monadic second-order logic. The claim follows from Courcelle’s theorem [14]. $\square$

Consider a plane graph G . If all vertices lie on the outerface, then G is 1-outerplanar. Otherwise, G is k -outerplanar if the removal of all vertices of the outerface results in a $(k - 1)$ -outerplanar graph. A graph is k -outerplanar if it has a k -outerplanar embedding in the plane [4].

Theorem 3 ([7]). *The treewidth of a k -outerplanar graph is at most $3k - 1$.*

Theorem 4. *There exists a polynomial-time approximation scheme for a maximum independent set of edges in planar graphs.*

We use Baker’s technique [4] in the proof of this theorem and we moved the proof to Appendix A.

Remark 3. It is easy to see that the treewidth algorithm, for graphs of treewidth k , can be implemented to run in $O(2^{O(k)}n)$ time. Since the treewidth of planar graphs is bounded by $O(\sqrt{n})$, this gives an exact algorithm for a maximum independent set of edges which runs in $O(2^{O(\sqrt{n})})$ time.

Remark 4. Blanchette, Kim and Vetta prove in [6] that there is a PTAS for edge-clique cover of planar graphs.

4.1 Planar graphs without triangle separator

A triangle separator (of a connected graph) is a triangle the removal of which disconnect the graph. We compute $\alpha'(G)$ for planar graphs G without triangle separator in the following theorem.

Theorem 5. *There exists an $O(n^{3/2})$ algorithm that computes a maximum independent set of edges in planar graphs without triangle separator.*

Proof. Let G be an embedding of a planar graph without triangle separator. We may assume that $G \not\cong K_4$. In all the faces that have length more than three add a new vertex and make it adjacent to all vertices in the face. Let G' be this graph. Give the edges of G a weight 1 and give the new edges a weight 0.

Let H be the dual of G' . In H the weight of an edge is the weight of the edge in G' that it crosses. We claim that a solution is obtained by computing a maximum weight matching in H [16].

Let M^d be a matching in H . We may assume that M^d contains no edges of weight 0. Then every edge of M^d crosses an edge of G . We claim that this is an independent set M of edges in G . Since M^d is a matching no two triangular faces of G' incident with different edges of M^d coincide. Assume that two edges of M lie in a triangle T of G . Then T cannot be a face of G' since M^d is a matching. But then T is a separator which contradicts our assumption.

Let M be an independent set of edges in G and let M^d be the corresponding edges of H . Since M is an independent set of edges no two edges of M lie in a triangle. Assume that M^d is not a matching, and let f be a common face of G' of two edges in M^d . Then f is contained in a face of G which is not a triangle. But then the edges of M are the same. $\square$

5 Rankwidth of edge-clique graphs of cocktail parties

In this section we show that the related problem to compute $\theta_e(G)$ is probably much harder than the independence number. We show that $\theta_e(G)$ relates to well-known open problems for cocktail party graphs G . Cocktail parties are a proper subclass of cographs.

Definition 2. *The cocktail party graph $cp(n)$ is the complement of a matching with $2n$ vertices.*

Gyárfás [21] showed an interesting lowerbound in the following theorem. Here, two vertices x and y are *equivalent* if they are adjacent and have the same closed neighborhood.

Theorem 6. *If a graph G has n vertices and contains neither isolated nor equivalent vertices then $\theta_e(G) \geq \log_2(n+1)$.*

Notice that a cocktail party graph has no equivalent vertices. Thus, by Theorem 6,

$$\theta_e(cp(n)) \geq \log_2(2n+1).$$

For the cocktail party graph an exact formula for $\theta_e(cp(n))$ appears in [19]. In that paper Gregory and Pullman (see also [17,27]) prove that

$$\lim_{n \rightarrow \infty} \frac{\theta_e(cp(n))}{\log_2(n)} = 1.$$

For a graph G we denote the vertex-clique cover number of G by $\kappa(G)$. Notice that, for a graph G ,

$$\theta_e(G) = \kappa(K_e(G)).$$

Albertson and Collins mention the following result (due to Shearer) [1] for the graphs $K_e^r(cp(n))$, defined recursively by $K_e^r(cp(n)) = K_e(K_e^{r-1}(cp(n)))$.

$$\alpha(K_e^r(cp(n))) \leq 3 \cdot (2^r)!$$

Thus, for $r = 1$, $\alpha(K_e(cp(n))) \leq 6$. However, the following is easily checked (see Lemma 3).

Lemma 6. For $n \geq 2$

$$\alpha(K_e(\text{cp}(n))) = 4.$$

Definition 3. A class of graphs $\mathcal{G}$ is χ -bounded if there exists a function f such that for every graph $G \in \mathcal{G}$,

$$\chi(G) \leq f(\omega(G)).$$

Dvořák and Král' prove that the class of graphs with rankwidth at most k is χ -bounded [15].

Theorem 7. The class of edge-clique graphs of cocktail parties has unbounded rankwidth.

Proof. It is easy to see that the rankwidth of any graph is at most one more than the rankwidth of its complement [34]. Assume that there is a constant k such that the rankwidth of $K_e(G)$ is at most k whenever G is a cocktail party graph. Let

$$\mathcal{K} = \{ \overline{K_e(G)} \mid G \simeq \text{cp}(n), n \in \mathbb{N} \}.$$

Then the rankwidth of graphs in $\mathcal{K}$ is uniformly bounded by $k + 1$. By the result of Dvořák and Král', there exists a function f such that

$$\log_2(2n + 1) \leq \theta_e(G) = \kappa(K_e(G)) \leq f(\alpha(K_e(G))) = 4f$$

for every cocktail party graph G . This contradicts Lemma 6 and Theorem 6. $\square$

6 Conclusion and discussions

In this paper we show polynomial-time algorithms and NP-completeness proofs for the independence number of edge-clique graphs of some classes of graphs. In particular we show that for cographs and distance-hereditary graphs the independence number of edge-clique graphs satisfies Formula (2). The results lead us ask new questions that for which other classes of graphs the formula also holds true and under what conditions the formula is true. For edge-clique cover problem we make a conjecture as follows.

Conjecture 1. The edge-clique cover problem is NP-complete for cographs.

We move the motivation of this conjecture to Appendix B.

Finally, trivially perfect graphs is the subclass of cographs that are chordal. A graph is trivially perfect if it does not contain C_4 nor P_4 as an induced subgraph. For this class of graphs we have equality:

Theorem 8. If a graph G is connected and trivially perfect then $\alpha'(G) = \theta_e(G)$.

Proof. If G contains only one vertex then $\alpha'(G) = \theta_e(G) = 0$. Assume that G contains more than one vertex. Since G is connected then $\alpha'(G) \geq \alpha(G)$. When a graph G is trivially perfect then the independence number is equal to the number of maximal cliques in G . Thus $\alpha(G) = \theta_e(G) \geq \alpha'(G)$. Finally, we conclude $\alpha'(G) = \theta_e(G)$. $\square$

References

1. Albertson, M. and K. Collins, Duality and perfection for edges in cliques, *Journal of Combinatorial Theory, Series B* **36** (1984), pp. 298–309.
2. Alcón, L., L. Faria, C. de Figueiredo and M. Gutierrez, The complexity of clique graph recognition, *Theoretical Computer Science* **410** (2009), pp. 2072–2083.
3. Anand, P., H. Escudero, R. Gera, S. Hartke and D. Stolee, On the hardness of recognizing triangular line graphs. *Discrete Mathematics* **312** (2012), pp. 2627–2638.
4. Baker, B., Approximation algorithms for NP-complete problems on planar graphs, *Journal of the ACM* **41** (1994), pp. 153–180.
5. Bandelt, H. and H. Mulder, Distance-hereditary graphs, *Journal of Combinatorial Theory, Series B* **41** (1986), pp. 182–208.
6. Blanchette, M. E. Kim and A. Vetta, Clique cover on sparse networks, *Proceedings ALENEX, SIAM* (2012), pp. 93–102.
7. Bodlaender, H., A partial k-arboretum of graphs with bounded treewidth, *Theoretical Computer Science* **209** (1998), pp. 1–45.
8. Calamoneri, T. and R. Petreschi, Edge-clique graphs and the λ -coloring problem, *J. Braz. Comput. Soc.* **7** (2001), pp. 38–47.
9. Cerioli, M., Clique graphs and edge-clique graphs, *Electronic Notes in Discrete Mathematics* **13** (2003), pp. 34–37.
10. Cerioli, M. and J. Szwarcfiter, A characterization of edge clique graphs, *Ars Combinatoria* **60** (2001), pp. 287–292.
11. Cerioli, M. and J. Szwarcfiter, Edge clique graphs and some classes of chordal graphs, *Discrete Mathematics* **242** (2002), pp. 31–39.
12. Chartrand, G. S. F. Kapoor, T. A. McKee and F. Saba, Edge-clique graphs, *Graphs and Combinatorics* **7** (1991), pp. 253–264.
13. Corneil, D., Y. Perl and L. Stewart, A linear recognition algorithm for cographs, *SIAM Journal on Computing* **14** (1985), pp. 926–934.
14. Courcelle, B., The monadic second-order logic of graphs. I. Recognizable sets of finite graphs, *Information and Computation* **85** (1990), pp. 12–75.
15. Dvořák, Z. and D. Král', Classes of graphs with small rank decompositions are χ -bounded. *European Journal of Combinatorics* **33** (2012), pp. 679–683.
16. Edmonds, J., Paths, trees, and flowers, *Canadian Journal of Mathematics* **17** (1965), pp. 449–467.
17. Gargano, L., J. Körner and U. Vaccaro, Sperner capacities, *Graphs and Combinatorics* **9** (1993), pp. 31–46.
18. Garey, M., and D. Johnson, The rectilinear Steiner tree problem is NP-complete, *SIAM J. Appl. Math.* **32** (1977), pp. 826–834.
19. Gregory, D. A. and N. J. Pullman, On a clique covering problem of Orlin, *Discrete Mathematics* **41** (1982), pp. 97–99.
20. Grötschel, M., L. Lovász and A. Schrijver, Polynomial algorithms for perfect graphs. In (Berge, Chvátal eds.): *Topics on perfect graphs*, North-Holland Mathematics Studies **88** (1984), pp. 325–356.
21. Gyárfás, A., A simple lower bound on edge covering by cliques, *Discrete Mathematics* **85** (1990), pp. 103–104.
22. Haxell, P., A. Kostochka and S. Thomassé, A stability theorem on fractional covering of triangles by edges, *European Journal of Combinatorics* **33** (2012), pp. 799–806.
23. Holyer, I., The NP-completeness of some edge-partition problems, *SIAM Journal on Computing* **4** (1981), pp. 713–717.

24. Hoover, D. N., Complexity of graph covering problems for graphs of low degree, *JCMCC* **11** (1992), pp. 187–208.
25. Howorka, E., A characterization of distance-hereditary graphs, *The Quarterly Journal of Mathematics* **28** (1977), pp. 417–420.
26. Kong, J. and Y. Wu, On economical set representations of graphs, *Discrete Mathematics and Theoretical Computer Science* **11** (2009), pp. 71–96.
27. Körner, J., Intersection number and capacities of graphs, *Discrete Mathematics* **142** (1995), pp. 169–184.
28. Kou, L. T., L. J. Stockmeyer and C. K. Wong, Covering edges by cliques with regard to keyword conflicts and intersection graphs, *Communications of the ACM* **21** (1978), pp. 135–139.
29. McKee, T. A., Clique multigraphs, In (Y. Alavi, F. R. K. Chung, R. L. Graham, D. F. Hsu eds.) *Second International Conference in Graph Theory, Combinatorics, Algorithms and Applications*, SIAM (1991), pp. 371–379.
30. Lakshmanan, S., C. Bujtás and Z. Tuza, Small edge sets meeting all triangles of a graph, *Graphs and Combinatorics* **28** (2012), pp. 381–392.
31. Lakshmanan, S. and A. Vijayakumar, Clique irreducibility of some iterative classes of graphs, *Discussiones Mathematicae Graph Theory* **28** (2008), pp. 307–321.
32. Le, V. B., Gallai graphs and anti-Gallai graphs, *Discrete Mathematics* **159** (1996), pp. 179–189.
33. Orlin, J., Contentment in graph theory: covering graphs with cliques, *Proceedings of the Nederlandse Academie van Wetenschappen, Amsterdam, Series A* **80** (1977), pp. 406–424.
34. Oum, S., *Graphs of bounded rankwidth*. PhD thesis, Princeton University, 2005.
35. Poljak, S., A note on stable sets and coloring of graphs, *Commentationes Mathematicae Universitatis Carolinae* **15** (1974), pp. 307–309.
36. Prisner, E., Graph dynamics, *Pitman Research Notes in Mathematics Series* Longman, Essex, (1995).
37. Pullman, N. J., Clique covering of graphs IV. Algorithms, *SIAM Journal on Computing* **13** (1984), pp. 57–75.
38. Raychaudhuri, A., Intersection number and edge clique graphs of chordal and strongly chordal graphs, *Congressus Numerantium* **67** (1988), pp. 197–204.
39. Raychaudhuri, A., Edge clique graphs of some important classes of graphs, *Ars Combinatoria* **32** (1991), pp. 269–278.
40. Scheinerman, E. and A. Trenk, On the fractional intersection number of a graph, *Graphs and Combinatorics* **15** (1999), pp. 341–351.
41. Sbihi, N., Algorithme de recherche d'un stable de cardinalité maximum dans un graphe sans étoile, *Discrete Mathematics* **29** (1980), pp. 53–76.

A Proof of Theorem 4

Theorem 9. *There exists a polynomial-time approximation scheme for a maximum independent set of edges in planar graphs.*

Proof. We use Baker’s technique [4]. Consider a plane embedding of G . Let L_0 be the set of vertices in the outerface. Remove the vertices of L_0 from G and let L_1 be the new outerface. Continuing this process partitions the vertices into layers.

Fix an integer k . For $i \in \{0, \dots, k-1\}$ and $j \in \mathbb{N} \cup \{-1, 0\}$, let G_{ij} be the subgraph of G induced by the vertices in layers L_t where

$$t \in \{jk + i + 1, jk + i + 2, \dots, jk + i + (k - 1)\} \text{ and } t \geq 0.$$

Notice that the outerplanarity of each G_{ij} is at most k . By lemma 5 there exists a linear-time algorithm which computes a maximum independent set of edges A_{ij} in G_{ij} .

Every edge in G connects two vertices in either the same layer or in two adjacent layers. For fixed i , the consecutive graphs G_{ij} skip two layers. So $\cup_j A_{ij}$ is an independent set of edges in G .

Let A be a maximum independent set of edges in G . Sum over all graphs G_{ij} the edges of A that are contained in G_{ij} . Then every edge of A is counted $k - 2$ times. Also, since A_{ij} is an exact solution for G_{ij} this is at least as big as the number of edges induced by A in G_{ij} . Therefore, the sum of A_{ij} is at least $k - 2$ times the optimum. If we take the maximum over $i \in \{0, \dots, k - 1\}$, we find an approximation of size at least $1 - \frac{2}{k}$ times the optimum.

This proves the theorem. $\square$

B Conjecture on edge-clique cover problem for cographs

According to Theorem 6 Gyárfás [21] result implies that the edge-clique cover problem is fixed-parameter tractable (see also [46]).

Cygan et al [44] show that, under the assumption of the exponential time hypothesis, there is no polynomial-time algorithm which reduces the parameterized problem $(\theta_e(G), k)$ to a kernel of size bounded by $2^{o(k)}$. In their proof the authors make use of the fact that $\theta_e(\text{cp}(2^\ell))$ is a [sic] “hard instance for the edge-clique cover problem, at least from a point of view of the currently known algorithms.” Note that, in contrast, the parameterized edge-clique partition problem can be reduced to a kernel with at most k^2 vertices [48]. (Mujuni and Rosamond also mention that the edge-clique cover problem probably has no polynomial kernel.) These observations lead us to investigate edge-clique graphs of cocktail parties in Section 5.

Let K_n^m denote the complete multipartite graph with m partite sets each having n vertices. Obviously, K_n^m is a cograph with $n \cdot m$ vertices.

Theorem 10 ([49]). *Assume that*

$$3 \leq m \leq n + 1.$$

Then $\theta_e(K_n^m) = n^2$ if and only if there exists a collection of at least $m - 2$ pairwise orthogonal Latin squares of order n .

Notice that, if there exists an edge-clique cover of K_n^m with n^2 cliques, then these cliques are mutually edge-disjoint. Finding the maximal number of mutually orthogonal Latin squares of order n is a renowned open problem. The problem has a wide field of applications, eg in combinatorics, designs of experiments, group theory and quantum informatics.

Unless n is a prime power, the maximal number of MOLS is known for only a few orders. We briefly mention a few results. Let $f(n)$ denote the maximal number of MOLS of order n . The well-known ‘Euler-spoiler’ shows that $f(n) = 1$ only for $n = 2$ and $n = 6$. Also, $f(n) \leq n - 1$ for all $n > 1$, and Chowla, Erdős and Straus [43] show that

$$\lim_{n \rightarrow \infty} f(n) = \infty.$$

Define

$$n_r = \max \{ n \mid f(n) < r \}.$$

A lowerbound for the speed at which $f(n)$ grows was obtained by Wilson, who showed that $n_r < r^{17}$ when r is sufficiently large [50]. Better bounds for n_r , for some specific values of r , were obtained by various authors (see eg [42]).

See eg [45] for some recent computational attempts to find orthogonal Latin squares. The problem seems extremely hard, both from a combinatorial and from a computational point of view [27]. Despite many efforts, the existence of three pairwise orthogonal Latin squares of order 10 is, as far as we know, still unclear.

Finally, the observations mentioned above lead us to conjecture that the edge-clique cover problem is NP-complete for cographs.

C Planar graphs of bounded treewidth

Theorem 11. *Let G be a planar graph with treewidth at most k . Then $\alpha'(G)$ can be computed in $O(2^{O(k)}n)$ time, where n is the number of vertices of G .*

Proof. Let $\{\mathcal{X}, T\}$ be a nice tree decomposition of $G = (V, E)$, where $\mathcal{X} = \{X_1, \dots, X_t\}$ is a family of subsets of V and T is a tree whose nodes are the subsets X_i [47]. Since $\{\mathcal{X}, T\}$ is nice, T is a rooted tree and there are four types of node i . We choose an arbitrary node r in T as the root of T .

For each node i , $G(i)$ is a subgraph of G formed by all nodes in sets X_j , with $j = i$ or j a descendant of i . Let $E(X_i)$ denote a set of the edges induced by X_i . For node i in tree decomposition and a subset $F \subseteq E(X_i)$, we define $\alpha'(i, F)$ as the

maximum cardinality of independent set A of edges of $G(i)$ with $A \cap E(X_i) = F$. Note that $\alpha'(i, F) = -\infty$ if A does not exist.

In the following, we consider the four types of nodes in T and calculate $\alpha'(i, F)$.

LEAF: Let i be a leaf node with $|X_i| = 1$. Then $F = \emptyset$ and $\alpha'(i, F) = 0$.

JOIN: Let i be a node with children j_1, j_2 . Then

$$\alpha'(i, F) = \alpha'(j_1, F) + \alpha'(j_2, F) - |F|. \quad (8)$$

INTRODUCE: Let i be a node with child node j such that $X_i = X_j \cup \{v\}$, for some vertex $v \in V$. Let $F \subseteq E(X_j)$. Then $\alpha'(i, F) = \alpha'(j, F)$. Let F' be a subset of the edges in $E(X_i)$ that are incident with v . Then if $F \cup F'$ is independent,

$$\alpha'(i, F \cup F') = \alpha'(j, F) + |F'|. \quad (9)$$

Otherwise, if F is not independent then $\alpha'(i, F \cup F') = -\infty$.

FORGET: Let i be a node with child node j such that $X_i = X_j \setminus \{v\}$, for some vertex $v \in V$. Let $F \subseteq E(X_i)$ and let F' be a subset of the edges in $E(X_j)$ that are incident with v . Then

$$\alpha'(i, F) = \max \{ \alpha'(j, F), \alpha'(j, F \cup F') \}. \quad (10)$$

For node i in tree decomposition we compute a table with all values $\alpha'(i, F)$ via dynamic programming. Since G is planar, each node i contains $O(k)$ edges. Thus for any type of nodes, a table contains $O(2^{O(k)})$ entries and can be calculated in $O(2^{O(k)})$ time. Therefore, $\alpha'(G) = \max \{ \alpha'(r, F) \}$ can be obtained in $O(2^{O(k)}n)$ time.

This completes the proof. $\square$

D The bibliography for the appendix

References

42. Brouwer, A. and G. Rees, More mutually orthogonal Latin squares, *Discrete Mathematics* **39** (1982), pp. 263–281.
43. Chowla, S., P. Erdős and E. G. Straus, On the maximal number of pairwise orthogonal Latin squares of a given order, *Canadian Journal of Mathematics* **12** (1960), pp. 204–208.
44. Cygan, M., M. Pilipczuk and M. Pilipczuk, Known algorithms for edge clique cover are probably optimal. Manuscript on ArXiv: 1203.1754v1, 2012.
45. Ma, F. and Zhang, J., Finding orthogonal Latin squares using finite model searching tools. Manuscript. To appear in *Science China – Information Sciences* 2011.
46. Gramm, J., J. Guo, F. Hüffner and R. Niedermeier, Data reduction, exact, and heuristic algorithms for clique cover, *Proceedings ALENEX'06*, SIAM (2006), pp. 86–94.
47. Kloks, T., *Treewidth – Computations and Approximations*, Springer, LNCS 842, 1994.
48. Mujuni, E. and F. Rosamond, Parameterized complexity of the clique partition problem, (J. Harland and P. Manyem, eds.) *Proceedings CATS'08*, ACS, CRPIT series **77** (2008), pp. 75–78.

49. Park, B., S. Kim and Y. Sano, The competition numbers of complete multipartite graphs and mutually orthogonal Latin squares, *Discrete Mathematics* **309** (2009), pp. 6464–6469.
50. Wilson, R., Concerning the number of mutually orthogonal Latin squares, *Discrete Mathematics* **9** (1974), pp. 181–198.